

Production of the neutral top-pion π_t^0 in association with a high- p_T jet at the LHC

Shi-Hai Zhu, Chong-Xing Yue, Wei Liu, and Li Ding

Department of Physics, Liaoning Normal University, Dalian 116029, P. R. China *

December 8, 2008

Abstract

In the framework of the topcolor-assisted technicolor ($TC2$) model, we study production of the neutral top-pion π_t^0 in association with a high- p_T jet at the LHC , which proceeds via the partonic processes $gg \rightarrow \pi_t^0 g$, $gq \rightarrow \pi_t^0 q$, $q\bar{q} \rightarrow \pi_t^0 g$, $gb(\bar{b}) \rightarrow \pi_t^0 b(\bar{b})$, and $b\bar{b} \rightarrow \pi_t^0 g$. We find that it is very challenging to detect the neutral top-pion π_t^0 via the process $pp \rightarrow \pi_t^0 + jet + X \rightarrow t\bar{t} + jet + X$, while the possible signatures of π_t^0 might be detected via the process $pp \rightarrow \pi_t^0 + jet + X \rightarrow (\bar{t}c + t\bar{c}) + jet + X$ at the LHC .

*E-mail: cxyue@lnnu.edu.cn

1. Introduction

The Higgs mechanism for the electroweak symmetry breaking (*EW**SB*) is still the untested part of the standard model (*SM*). Searching for the *SM* Higgs boson is one of the main tasks of the forthcoming Large Hadron Collider (*LHC*), which has considerable capability to discover and measure almost all of its quantum properties [1]. However, if the *LHC* finds evidence for a new scalar state, it may not necessarily be the *SM* Higgs boson. Most of new physics models beyond the *SM* predict the existence of new scalar states. These new particles may have production cross sections and branching ratios which differ from those of the *SM* Higgs boson. Distinguishing the various new physics scenarios is an important task for current and near future high energy collider experiments. Thus, studying the production and decay of the new scalar states at the *LHC* is of special interest.

Due to the large gluon luminosity, the main production mechanism for a scalar Higgs boson at the *LHC* is the partonic gluon fusion process $gg \rightarrow H$ [2], which is the so-called inclusive single Higgs boson production channel. In order to fully explore the Higgs detection capabilities of the *LHC*, one should investigate more exclusive channels, like e.g. Higgs production in association with a high- p_T hadronic jet [3]. The main advantage of this channel is the richer kinematical structure of the events which allows for refined cuts increasing the signal-to-background ratio. So far, this production channel has been extensively studied in the *SM* [4,5]. In the minimal supersymmetric standard model (*MSSM*), the analogous process, i.e. scalar Higgs production in association with a high- p_T jet was also extensively studied in Refs.[6,7].

Among various kinds of dynamical *EW**SB* theories, the topcolor scenario is attractive because it can explain the large top quark mass and provides a possible *EW**SB* mechanism [8]. The topcolor-assisted technicolor (*TC*2) model [9] is one of the phenomenologically viable models, which has all essential features of the topcolor scenario. This model predicts three *CP* odd top-pions ($\pi_t^0, \pi_t^\pm$) with large Yukawa couplings to the third family. The aim of this paper is to consider the production of the neutral top-pion π_t^0 associated with a high- p_T jet and compare our results with those for the Higgs boson from the *SM* or the

MSSM. We hope that our work can help the upcoming *LHC* to test topcolor scenario and to differentiate various kinds of new physics models.

In the rest of this paper, we will give our results in detail. In section 2, we will calculate the production cross section of the hadronic process $pp \longrightarrow \pi_t^0 + jet + X$ and give a simply phenomenological analysis at the *LHC*. Our conclusion is represented in section 3.

2. Production of the neutral top-pion π_t^0 associated with a high- p_T jet

In the *TC2* model [9], topcolor interactions, which are not flavor-universal and mainly couple to third generation fermions, generally generate small contributions to *EW**SB* and give rise to the main part of the top quark mass. Thus, the top-pions $\pi_t^{0,\pm}$ have large Yukawa couplings to the third generation fermions. Such features can result in large tree-level flavor changing couplings of the top-pions to the fermions when one writes the interactions in the fermion mass eigen-basis. Just as for the *SM* Higgs boson, the couplings of the top-pion to a pair of quarks are proportion to the quark masses. The explicit form for the couplings of the neutral top-pion π_t^0 to quarks, which are related to our calculation, can be written as [9,10]:

$$\frac{im_t}{\sqrt{2}F_t} \frac{\sqrt{\nu_W^2 - F_t^2}}{\nu_W} [k_{UR}^{tt} k_{UL}^{t*} \bar{t} \gamma^5 t \pi_t^0 + \frac{m_b - m'_b}{m_t} \bar{b} \gamma^5 b \pi_t^0 + k_{UR}^{tc*} k_{UL}^{tt} \bar{t} P_R c \pi_t^0], \quad (1)$$

where $\nu_W = \nu/\sqrt{2} \approx 174 GeV$, $P_R = (1 + \gamma^5)/2$ is the right-handed projection operator, $F_t \approx 50 GeV$ is the top-pion decay constant, and $m'_b \approx 0.1\epsilon m_t$ is the part of the bottom quark mass generated by extended technicolor interactions. $k_{UL(R)}$ are rotation matrices that diagonalize the up-quark mass matrix M_U for which the Cabibbo-Kobayashi-Maskawa (*CKM*) matrix is defined as $V_{CKM} = k_{UL}^+ k_{DL}$. To yield a realistic form of V_{CKM} , it has been shown that the values of the matrix elements $k_{UL(R)}^{ij}$ can be taken as [10]:

$$k_{UL}^{tt} \approx 1, \quad k_{UR}^{tt} = 1 - \epsilon, \quad k_{UR}^{tc} \leq \sqrt{2\epsilon - \epsilon^2}. \quad (2)$$

In our numerical estimation, we will take $k_{UR}^{tc} = \sqrt{2\epsilon - \epsilon^2}$ and take ϵ as a free parameter, which is assumed to be in the range of $0.01 \sim 0.1$.




Figure 1: Feynman diagrams for the partonic processes contributing to the hadronic process $pp \rightarrow \pi_t^0 + jet + X$ at the leading order. The other diagrams obtained by exchanging the gluons or exchanging π_t^0 are not shown here.

Similar to the Higgs boson predicted by the *SM* or the *MSSM*, the neutral top-pion π_t^0 can be produced at the *LHC* in association with a high- p_T jet through three partonic processes: gluon fusion ($gg \longrightarrow g\pi_t^0$), quark-gluon scattering ($q(\bar{q})g \longrightarrow q(\bar{q})\pi_t^0$), and quark-antiquark annihilation ($q\bar{q} \longrightarrow g\pi_t^0$). Although the gluon fusion and quark-gluon scattering partonic processes give main contributions to the production cross section for the hadronic process $pp \longrightarrow \pi_t^0 + jet + X$ at the *LHC*, our numerical analysis include all of above three processes, which proceed at one-loop, as shown in *Fig.1(a) ~ (e)*. Considering the small value of the decay constant F_t and the relatively large bottom-quark mass, we also consider the contributions of the tree-level partonic processes $gb \longrightarrow \pi_t^0 b$ and $b\bar{b} \longrightarrow \pi_t^0 g$, as shown in *Fig.1(f) ~ (h)*.

The variant amplitudes corresponding to the Feynman diagrams as shown in *Fig.1* can be written as:

$$M_{(a)} = \frac{i}{4\sqrt{2}\pi^2} T_{ij}^{c_3} T_{ji}^{c_4} f^{c_1 c_2 c_3} g_s^3 \frac{m_t^2 (1 - \varepsilon) \sqrt{\nu_W^2 - F_t^2}}{F_t \nu_W (p_1 + p_2)^2} C_{0(a)} [(p_2 - p_1)_\mu \epsilon(p_1) \cdot \epsilon(p_2) + (-2p_2 - p_1) \cdot \epsilon(p_1) \epsilon_\mu(p_2) + (2p_1 + p_2) \cdot \epsilon(p_2) \epsilon_\mu(p_1)] \epsilon^{\mu\nu\rho\sigma} p_{3\rho} p_{4\sigma} \epsilon_\nu(p_3), \quad (3)$$

$$M_{(b)} = \frac{i}{4\sqrt{2}\pi^2} T_{ij}^{c_3} T_{ji}^{c_4} f^{c_1 c_2 c_3} g_s^3 \frac{m_t^2 (1 - \varepsilon) \sqrt{\nu_W^2 - F_t^2}}{F_t \nu_W (p_1 - p_3)^2} C_{0(b)} [(-p_2 - p_1) \cdot \epsilon(p_3) \epsilon_\mu(p_1) + (p_2 - p_3) \cdot \epsilon(p_1) \epsilon_\mu(p_3) + (p_3 + p_1)_\mu \epsilon(p_1) \cdot \epsilon(p_3)] \epsilon^{\mu\nu\rho\sigma} p_{4\rho} (p_1 - p_3)_\sigma \epsilon_\nu(p_2), \quad (4)$$

$$M_{(c)} = \frac{i}{4\sqrt{2}\pi^2} T_{ij}^{c_1} T_{jk}^{c_2} T_{ki}^{c_3} g_s^3 \frac{m_t (1 - \varepsilon) \sqrt{\nu_W^2 - F_t^2}}{F_t \nu_W} \\ (-D_{0(c)} m_t^3 \epsilon^{\mu\nu\rho\sigma} p_{1\sigma} + D_{0(c)} m_t^3 \epsilon^{\mu\nu\rho\sigma} p_{4\sigma} - D_{0(c)} m_t \epsilon^{\rho\sigma\alpha\beta} p_{1\sigma} p_{2\alpha} p_{4\beta} g^{\mu\nu} \\ + D_{0(c)} m_t \epsilon^{\nu\sigma\alpha\beta} p_{1\sigma} p_{2\alpha} p_{4\beta} g^{\mu\rho} - D_{0(c)} m_t \epsilon^{\nu\rho\sigma\alpha} p_1^\mu p_{2\sigma} p_{4\alpha} + D_{0(c)} m_t \epsilon^{\nu\rho\sigma\alpha} p_{1\sigma} p_{4\alpha} p_2^\mu \\ - D_{0(c)} m_t \epsilon^{\nu\rho\sigma\alpha} p_{1\sigma} p_{2\alpha} p_4^\mu - D_{0(c)} m_t \epsilon^{\mu\sigma\alpha\beta} p_{1\sigma} p_{2\alpha} p_{4\beta} g^{\nu\rho} + 2D_{(c)}^\nu m_t \epsilon^{\mu\rho\sigma\alpha} p_{1\sigma} p_{4\alpha} \\ + 2D_{(c)}^\nu m_t \epsilon^{\mu\rho\sigma\alpha} p_{2\sigma} p_{4\alpha} + D_{0(c)} m_t \epsilon^{\mu\rho\sigma\alpha} p_{2\sigma} p_{4\alpha} p_1^\nu - D_{0(c)} m_t \epsilon^{\mu\rho\sigma\alpha} p_{1\sigma} p_{4\alpha} p_2^\nu \\ + D_{0(c)} m_t \epsilon^{\mu\rho\sigma\alpha} p_{1\sigma} p_{2\alpha} p_4^\nu - 2D_{(c)}^\nu m_t \epsilon^{\mu\rho\sigma\alpha} p_{2\sigma} p_{4\alpha} - D_{0(c)} m_t \epsilon^{\mu\nu\sigma\alpha} p_{2\sigma} p_{4\alpha} p_1^\rho \\ - D_{0(c)} m_t \epsilon^{\mu\nu\sigma\alpha} p_{1\sigma} p_{4\alpha} p_2^\rho + D_{0(c)} m_t \epsilon^{\mu\nu\sigma\alpha} p_{1\sigma} p_{2\alpha} p_4^\rho + D_{\nu(c)} D_{(c)}^\nu m_t \epsilon^{\mu\nu\rho\sigma} p_{1\sigma} \\ - D_{\nu(c)} D_{(c)}^\nu m_t \epsilon^{\mu\nu\rho\sigma} p_{4\sigma} - 2D_{\nu(c)} \cdot p_2 m_t \epsilon^{\mu\nu\rho\sigma} p_{1\sigma} + 2D_{\nu(c)} \cdot p_2 m_t \epsilon^{\mu\nu\rho\sigma} p_{4\sigma} \\ - 2D_{\nu(c)} \cdot p_4 m_t \epsilon^{\mu\nu\rho\sigma} p_{2\sigma} + D_{0(c)} p_1 \cdot p_2 m_t \epsilon^{\mu\nu\rho\sigma} p_{4\sigma} - D_{0(c)} p_1 \cdot p_4 m_t \epsilon^{\mu\nu\rho\sigma} p_{2\sigma}$$

$$+D_{0(c)}p_2 \cdot p_2 m_t \epsilon^{\mu\nu\rho\sigma} p_{1\sigma} - D_{0(c)}p_2 \cdot p_4 m_t \epsilon^{\mu\nu\rho\sigma} p_{1\sigma} \epsilon_\mu(p_1) \epsilon_\nu(p_2) \epsilon_\rho(p_3) \quad (5)$$

$$M_{(d)} = \frac{1}{4\sqrt{2}\pi^2} T_{ij}^{c_1} T_{kl}^{c_1} T_{lk}^{c_2} g_s^3 \frac{m_t^2(1-\varepsilon)\sqrt{\nu_W^2 - F_t^2}}{F_t \nu_W (p_1 - p_3)^2} C_{0(d)} \bar{u}(p_3) \gamma^\mu u(p_1) g_{\mu\nu} \epsilon^{\nu\lambda\rho\sigma} p_{4\rho} (p_1 - p_3)_\sigma \epsilon_\lambda(p_2), \quad (6)$$

$$M_{(e)} = \frac{1}{4\sqrt{2}\pi^2} T_{ij}^{c_1} T_{kl}^{c_1} T_{lk}^{c_2} g_s^3 \frac{m_t^2(1-\varepsilon)\sqrt{\nu_W^2 - F_t^2}}{F_t \nu_W (p_1 + p_2)^2} C_{0(e)} \bar{u}(p_2) \gamma^\mu v(p_1) g_{\mu\nu} \epsilon^{\nu\lambda\rho\sigma} p_{3\rho} p_{4\sigma} \epsilon_\lambda(p_3), \quad (7)$$

$$M_{(f)} = \frac{1}{\sqrt{2}} T_{ij}^{c_1} g_s \frac{1}{(p_1 + p_2)^2 - m_b^2} \frac{m_b - m'_b}{F_t} \frac{\sqrt{\nu_W^2 - F_t^2}}{\nu_W} \bar{u}(p_3) \gamma^5 (\not{p}_1 + \not{p}_2 - m_b) \gamma^\mu u(p_1) \epsilon_\mu(p_2), \quad (8)$$

$$M_{(g)} = \frac{1}{\sqrt{2}} T_{ij}^{c_1} g_s \frac{1}{(p_1 - p_3)^2 - m_b^2} \frac{m_b - m'_b}{F_t} \frac{\sqrt{\nu_W^2 - F_t^2}}{\nu_W} \bar{u}(p_4) \gamma^\mu (\not{p}_1 - \not{p}_3 - m_b) \gamma^5 u(p_1) \epsilon_\mu(p_2), \quad (9)$$

$$M_{(h)} = \frac{1}{\sqrt{2}} T_{ij}^{c_1} g_s \frac{1}{(p_1 - p_3)^2 - m_b^2} \frac{m_b - m'_b}{F_t} \frac{\sqrt{\nu_W^2 - F_t^2}}{\nu_W} \bar{v}(p_2) \gamma^\mu (\not{p}_1 - \not{p}_3 - m_b) \gamma^5 u(p_1) \epsilon_\mu(p_4). \quad (10)$$

Here p_1, p_2 are the momenta of the incoming states, and p_3, p_4 are the momenta of the outgoing final states. The T_{ij}^c are the $SU(3)$ color matrices and the $f^{c_1 c_2 c_3}$ are the anti-symmetric $SU(3)$ structure constants in which i, j are the color indices and c_1, c_2, c_3 are the indices of gluon. The three-point and four-point standard functions C_0, D_0, D_1, D_ν [11,12] for different Feynman diagrams are defined as:

$$\begin{aligned} C_{0(a)} &= C_{0(a)}(p_1 + p_2, -p_3, m_t, m_t, m_t), C_{0(b)} = C_{0(b)}(p_1 - p_3, p_2, m_t, m_t, m_t), \\ C_{0(d)} &= C_{0(d)}(p_1 - p_3, p_2, m_t, m_t, m_t), C_{0(e)} = C_{0(e)}(p_1 + p_2, -p_3, m_t, m_t, m_t); \\ D_{0(c)} &= D_{0(c)}(p_1, p_2, -p_3, m_t, m_t, m_t, m_t), D_{1(c)} = D_{1(c)}(p_1, p_2, -p_3, m_t, m_t, m_t, m_t), \\ D_{\nu(c)} &= p_{1\nu} * D_{1(c)}(1) + p_{2\nu} * D_{1(c)}(2) + p_{3\nu} * D_{1(c)}(3). \end{aligned}$$

Each loop diagram is composed of some scalar loop functions, which are calculated by using LoopTools [12].

The hadronic cross section at the *LHC* is obtained by convoluting the partonic cross sections with the parton distribution functions (*PDFs*). In our numerical calculation, we

will use *CTEQ6L PDFs* [13] for the gluon and quark *PDFs*. The renormalization scale μ_R and the factorization scale μ_F are chosen to be $\mu_R = \mu_F = m_{\pi_t}$ for the gluons and the light quarks, and to be $\mu_R = \mu_F = m_{\pi_t}/4$ for the bottom-quark, in which m_{π_t} is the mass of the neutral top-pion π_t^0 . To make our predictions more realistic and high- p_T jet not too close to the beam axis, we require that the transverse momentum p_T and pseudorapidity η of the hadronic jet satisfy: $p_T > 30\text{GeV}$ and $|\eta| < 4.5$, which have been used in previous *MSSM* studies for the *LHC* [6,7].




Figure 2: The total cross section of the hadronic process $pp \rightarrow \pi_t^0 + jet + X$ as a function of the π_t^0 mass m_{π_t} for three values of the free parameter ϵ .

From the above discussions we can see that the production cross section σ for the hadronic process $pp \rightarrow \pi_t^0 + jet + X$ is dependent on the free parameters ϵ and m_{π_t} . Similar with Ref.[14], we will assume that the free parameters ϵ and m_{π_t} are in the range of $0.01 \sim 0.1$ and $200\text{GeV} \sim 500\text{GeV}$, respectively.

Our numerical results are shown in *Fig.2*, in which we plot the cross section σ as a function of the mass parameter m_{π_t} for three values of the parameter ϵ . One can see from *Fig.2* that σ is insensitive to the free parameter ϵ . For $\epsilon = 0.05$ and $200\text{GeV} \leq m_{\pi_t} \leq 500\text{GeV}$, the value of the production cross section σ is in the range of $18.3\text{pb} \sim 2.1\text{pb}$.

Observably, if we assume that the π_t^0 mass m_{π_t} is equal to that of the *SM* Higgs boson H or the *MSSM* Higgs boson H^0 , the cross section for the production of the neutral top-pion π_t^0 associated with a high- p_T is significantly larger than that of the *SM* Higgs boson H [4,5] or the *MSSM* Higgs boson H^0 [6,7]. This is because the $\pi_t^0 t \bar{t}$ coupling is larger than that for the *SM* Higgs boson H or the *MSSM* Higgs boson H^0 .




Figure 3: The hadronic cross sections for different partonic processes as function of the π_t^0 mass m_{π_t} for the free parameter $\varepsilon = 0.5$.

To see contributions of the different partonic processes to the total hadronic cross section, we plot the hadronic cross sections of the partonic processes $gg \rightarrow \pi_t^0 g$, $qg \rightarrow q\pi_t^0$ ($q = u, c, d, s, \bar{u}, \bar{c}, \bar{d}, \bar{s}$), $q\bar{q} \rightarrow \pi_t^0 g$ ($q = u, c, d, s$), $gb(\bar{b}) \rightarrow \pi_t^0 b(\bar{b})$, and $b\bar{b} \rightarrow \pi_t^0 g$ for $\varepsilon=0.05$ in *Fig.3*. We see that the production of the neutral top-pion π_t^0 in association with a high- p_T jet is dominated by the partonic process $gg \rightarrow \pi_t^0 g$, which is similar with the Higgs boson production associated with a high- p_T in the *SM* and the *MSSM*. However, for the *MSSM* model, the contributions of the $b\bar{b}$ channel can be significantly large, depending the free parameters. However, this is not the case for the *TC2* model. For $0.02 \leq \varepsilon \leq 0.08$ and $200\text{GeV} \leq m_{\pi_t} \leq 500\text{GeV}$, the hadronic cross section for the partonic process $b\bar{b} \rightarrow \pi_t^0 g$ is only in the range of $1.6\text{fb} \sim 46\text{fb}$, which is several orders

of magnitude smaller than that for the partonic process $gg \rightarrow \pi_t^0 g$.




Figure 4: The number of the $tc + jet$ event as a function of the π_t^0 mass m_{π_t} for three values of the parameter ϵ .

It is well known that the mass of the *SM* Higgs boson H is generally smaller than 200GeV , one can use the decay channels $H \rightarrow \gamma\gamma$, $H \rightarrow \tau^+\tau^-$ or $H \rightarrow W^+W^-$ to consider the *SM* Higgs boson signatures generated by the hadronic process $pp \rightarrow H + jet + X$ at the *LHC* [5]. For the neutral top-pion π_t^0 , its main decay modes are $t\bar{t}$, $\bar{t}c(t\bar{c})$, $b\bar{b}$, gg , and $\gamma\gamma$. For $m_t \leq m_{\pi_t} \leq 2m_t$, π_t^0 mainly decays to $\bar{t}c$ and $t\bar{c}$. It has been shown that the value of the branching ratio $Br(\pi_t^0 \rightarrow \bar{t}c + t\bar{c})$ is larger than 90% for $m_{\pi_t} = 250\text{GeV}$ and $\epsilon \geq 0.02$ [15]. Thus, for $m_t < m_{\pi_t} \leq 2m_t$, the production of neutral top-pion π_t^0 associated with a high- p_T hadronic jet can easily transfer to the $tc + jet$ event. This final state generates characteristic signatures at the *LHC* experiments. So we further calculate its production rate. We find that, for $\epsilon \leq 0.08$ and $m_{\pi_t} \leq 350\text{GeV}$, the production cross section of the hadronic process $pp \rightarrow (\bar{t}c + t\bar{c}) + jet + X$ is larger than 19.4pb . If we assume the yearly integrated luminosity $\mathcal{L}_{int} = 100\text{fb}^{-1}$ for the *LHC* with $\sqrt{s} = 14\text{TeV}$, then there will be $1.94 \times 10^6 \sim 5.3 \times 10^5$ $tc + jet$ events to be generated per year for $0.02 \leq \epsilon \leq 0.08$ and

$200\text{GeV} \leq m_{\pi_t} \leq 340\text{GeV}$, as shown in Fig.4.

For the $tc + jet$ event, the peak of the invariant mass distribution of tc is narrow. To identify tc , one needs reconstruct top quark from its mainly decay mode Wb and the b-tagging and c-tagging are also needed. Furthermore, in the case of the W hadronic decay, the $tc + jet$ event will generate the $bjjcj$ final state, while for the W leptonic decay, it will generate the $blvcj$ final state. For the former final state, the SM background is $jjjjj$ and the SM backgrounds of the later final state mainly come from the $t\bar{t}$, tW and $Wjjj$ production process, which have been analyzed in Ref.[16]. They have shown that suitable kinematical cuts on the observed particles is more than enough to obtain a clear and statistically meaningful flavor-changing signal. Thus we expect that the possible signatures of the neutral top-pion π_t^0 might be detected via the decay channel $\pi_t^0 \rightarrow \bar{t}c + t\bar{c}$ at the LHC experiments.

For $m_{\pi_t} > 2m_t$, the neutral top-pion π_t^0 mainly decays to $t\bar{t}$ and the hadronic process $pp \rightarrow \pi_t^0 + jet + X$ can give rise to the $t\bar{t} + jet$ event. Its production rate can reach 15pb for $m_{\pi_t} \geq 400\text{GeV}$ and $\varepsilon \leq 0.08$. This kind of events have been calculated at NLO in the SM [17]. It has shown that, for the renormalization and factorization scales having $\mu_R = \mu_F = \mu = m_t$, the NLO cross section for $t\bar{t} + jet$ production at the LHC is larger than 500pb. Thus, the production cross section of the $t\bar{t} + jet$ final state coming from $TC2$ is smaller than that coming from the SM by at least two orders of magnitude. It is very challenging to detect the possible signals of π_t^0 via the process $pp \rightarrow \pi_t^0 + jet + X \rightarrow t\bar{t} + jet + X$.

3. Conclusion

The production of a scalar state (the SM Higgs boson, the $MSSM$ Higgs boson, etc) associated with a high- p_T jet allows for refined cuts increasing the signal-to-background ratio, which is considered advantageous for scalar detection even though its production rate is lower than that for totally inclusive single scalar state production. In the context of the $TC2$ model, we consider the production of the neutral top-pion π_t^0 accompanied by a high- p_T jet at the LHC . This production channel proceeds by the partonic processes $gg \rightarrow \pi_t^0 g$, $gq \rightarrow \pi_t^0 q$, $q\bar{q} \rightarrow \pi_t^0 g$, $gb(\bar{b}) \rightarrow \pi_t^0 b(\bar{b})$, and $b\bar{b} \rightarrow \pi_t^0 g$. We find that,

for m_{π_t} equaling to the mass of the scalar state predicted by the *MSSM*, the hadronic production cross section of the process $pp \longrightarrow \pi_t^0 + jet + X$ is much larger than that for the *MSSM* scalar state. For $m_t < m_{\pi_t} \leq 2m_t$, the main decay channel is $\pi_t^0 \rightarrow \bar{t}c + t\bar{c}$. There will be a large number of the $tc + jet$ events to be generated which can generate characteristic signal at the *LHC* experiment. So we might detect the possible signatures of the neutral top-pion π_t^0 via the process $pp \longrightarrow \pi_t^0 + jet + X \rightarrow (\bar{t}c + t\bar{c}) + jet + X$ at the *LHC*.

Acknowledgments

Shi-Hai Zhu would like to thank Lei Wang for useful discussions. This work was supported in part by the National Natural Science Foundation of China under Grants No.10675057 and Foundation of Liaoning Educational Committee(2007T086).

References

- [1] For a recent review, see: N. E. Adam et al, *arXiv: 0803.1154*[hep-ph].
- [2] H. M. Georgi, S. L. Glashow, M. E. Machacek and D. V. Nanopoulos, *Phys. Rev. Lett.* **40**, 692(1978).
- [3] M. Chaichian, I. Liede, J. Lindfors, D. P. Roy, *Phys. Lett. B***198**, 416(1987)[Erratum-*ibid B***205**, 595(1988)]; R. K. Ellis, I. Hinchliffe, M. Soldate, J. J. van der Bij, *Nucl. Phys. B***297**, 221(1988); U. Baur, E. W. Nigel Glover, *Nucl. Phys. B***339**, 38(1990).
- [4] S. Abdellin et al., *Phys. Lett. B***431**, 410(1998); B. Mellado, W. Quayle, S. L. Wu, *Phys. Lett. B***611**, 60(2005); *Phys. Rev. D***76**, 093007(2007).
- [5] C. Anastasiou, K. Melnikov and F. Petriello, *Phys. Rev. Lett.* **93**, 262002(2004); *Nucl. Phys. B***724**, 197(2005); G. Bozzi, S. Catani, D. de Florian and M. Grazzini, *Phys. Lett. B* **564**, 65(2003); S. Catani, D. de Florian, M. Grazzini and P. Nason, *JHEP* **0307**, 028(2003); D. de Florian, M. Grazzini and Z. Kunszt, *Phys. Rev. Lett.* **82**, 5209(1999); D. de Florian, Z. Kunszt and W. Vogelsang, *JHEP* **0602**, 047(2006).
- [6] O. Brein and W. Hollik, *Phys. Rev. D***68**, 095006(2003); B. Field, S. Dawson and J. Smith, *Phys. Rev. D***69**, 074013(2004); U. Langenegger, M. Spira, A. Statradumov, P. Trucb, *JHEP* **0606**, 035(2006).
- [7] O. Brein and W. Hollik, *Phys. Rev. D***76**, 035002(2007); K. Bonciani, G. Degrassi, A. Vicini, *JHEP* **0711**, 095(2007); O. Brein and W. Hollik, *arXiv: 0710.4781*[hep-ph].
- [8] C. T. Hill and E. H. Simmons, *Phys. Rept.* **381**, 235(2003); [*Erratum-ibid*, **390**, 553(2004)].
- [9] C. T. Hill, *Phys. Lett. B* **345**, 483(1995); K. D. Lane and E. Eichten, *Phys. Lett. B* **352**, 382(1995); K. D. Lane, *Phys. Lett. B* **433**, 96(1998); G. Cvetcic, *Rev. Mod. Phys.* **71**, 513(1999).

- [10] G. Burdman, *Phys. Rev. Lett.* **83**, 2888(1999); H.-J. He, C.-P. Yuan, *Phys. Rev. Lett.* **83**, 28(1999); H.-J. He, S. Kanemura, C.-P. Yuan, *Phys. Rev. Lett.* **89**, 101803(2002).
- [11] G. 't Hooft and M. J. G. Veltman, *Nucl. Phys. B***153**, 365(1979).
- [12] T. Hahn, M. Perez-Victora, *Comput. Phys. Commun.* **118**, 153(1999); T. Hahn, *Nucl. Phys. Proc. Suppl.* **135**, 333(2004).
- [13] J. Pumplin et al. (CTEQ Collaboration), *JHEP* **02**, 032(2006).
- [14] C. -X. Yue, Z. -J. Zong, L. -L. Xu and J. -X. Chen, *Phys. Rev D***73**, 015006(2006).
- [15] C. -X. Yue, Y. -P. Kuang, and G. -G. Lu, *J. Phys. G***23**, 163(1997); C. -X. Yue, Q. -J. Xu, G. -L. Liu, and J. -T. Li, *Phys. Rev D***63**, 115002(2001).
- [16] F. del Aguila and J. A. Aguilar-Saavedra, *Nucl. Phys. B***576**, 56(2000).
- [17] S. Diffmaier, P. Uwer and S. Weinzierl, *Phys. Rev. Lett.* **98**, 262002(2007);*arXiv: 0804.4389*[hep-ph].